

## RESEARCH ARTICLE

## Open Access

# Functional genomics of the horn fly, *Haematobia irritans* (Linnaeus, 1758)

Lorena Torres<sup>1</sup>, Consuelo Almazán<sup>1\*</sup>, Nieves Ayllón<sup>2</sup>, Ruth C Galindo<sup>2</sup>, Rodrigo Rosario-Cruz<sup>3</sup>, Héctor Quiroz-Romero<sup>4</sup>, José de la Fuente<sup>2,5</sup>

## Abstract

**Background:** The horn fly, *Haematobia irritans* (Linnaeus, 1758) (Diptera: Muscidae) is one of the most important ectoparasites of pastured cattle. Horn flies infestations reduce cattle weight gain and milk production. Additionally, horn flies are mechanical vectors of different pathogens that cause disease in cattle. The aim of this study was to conduct a functional genomics study in female horn flies using Expressed Sequence Tags (EST) analysis and RNA interference (RNAi).

**Results:** A cDNA library was made from whole abdominal tissues collected from partially fed adult female horn flies. High quality horn fly ESTs (2,160) were sequenced and assembled into 992 unigenes (178 contigs and 814 singlets) representing molecular functions such as serine proteases, cell metabolism, mitochondrial function, transcription and translation, transport, chromatin structure, vitellogenesis, cytoskeleton, DNA replication, cell response to stress and infection, cell proliferation and cell-cell interactions, intracellular trafficking and secretion, and development. Functional analyses were conducted using RNAi for the first time in horn flies. Gene knockdown by RNAi resulted in higher horn fly mortality (protease inhibitor functional group), reduced oviposition (vitellogenin, ferritin and vATPase groups) or both (immune response and 5'-NUC groups) when compared to controls. Silencing of ubiquitination ESTs did not affect horn fly mortality and oviposition while gene knockdown in the ferritin and vATPase functional groups reduced mortality when compared to controls.

**Conclusions:** These results advanced the molecular characterization of this important ectoparasite and suggested candidate protective antigens for the development of vaccines for the control of horn fly infestations.

## Background

The horn fly, *Haematobia irritans* (Linnaeus, 1758) (Diptera: Muscidae) is one of the most important ectoparasites of pastured cattle [1]. This fly was originally introduced from Europe and currently represents a tremendous health problem for cattle in the Americas from Southern Canada to Argentina [2]. Although horn flies parasitize mainly cattle, occasionally they feed on horses, sheep and dogs [3].

The developmental cycle of *H. irritans* is very short, taking from 10 to 14 days to complete. Larvae and pupae develop on dung and once the flies emerge from pupae, immediately start and remain feeding on cattle

during their whole life. Flies leave the host only to move to others or to lay eggs on fresh manure [1]. Both males and females feed 24 to 38 times per day ingesting an average of 14.3 mg blood per fly [4].

Horn flies infestations interfere with animal feeding, thus producing significant reductions in weight gain and milk production [5,6]. The economic impact of *H. irritans* on livestock in the United States was estimated in approximately US\$1 billion annually [7,8]. In dairy cattle, infestations higher than 200 flies per animal produce a loss of 520 ml milk and 28 kg weight daily [6]. In beef cattle, *H. irritans* infestations can cause a reduction of 8.1 kg weight daily [5]. Moreover, the skin lesions caused by the intermittent feeding of horn flies produce significant hide damages, affecting considerably the leather industry [9]. Additionally, horn flies are mechanical vectors of different pathogens that cause disease in cattle [10-14].

\* Correspondence: [consuelo\\_almazan@yahoo.com.mx](mailto:consuelo_almazan@yahoo.com.mx)

<sup>1</sup>Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Tamaulipas, Km. 5 carretera Victoria-Mante, CP 87000 Ciudad Victoria, Tamaulipas, Mexico

Full list of author information is available at the end of the article

The control of horn flies has been primarily based on the use of chemical insecticides [15,16]. This control strategy has been partially successful but has resulted in the selection of flies resistant to most commercially available insecticides [15-17]. In addition to resistance, chemical insecticides affect other living organisms, contribute to environmental pollution and contaminate cattle products for human consumption.

Recently, research has been conducted to develop new horn fly control strategies that are cost-effective and environmentally friendly. The efficacy of the entomopathogenic fungi, *Metarhizium anisopaliniae*, against horn fly larvae was very high in vitro [18]. However, field application of entomopathogenic fungi for biological control of horn flies is difficult. The use of female-specific conditional lethality systems has been also considered but not yet developed [19].

The immunological control of ectoparasite infestations was demonstrated through cattle vaccination against tick infestations [20,21]. The effect of anti-tick vaccines on the reduction of cattle tick infestations and the transmission of some tick-borne pathogens [21-23] and preliminary results obtained in insect vector species [24-32] have provided evidence that protective antigens may be used for development of vaccines with the dual target control of both arthropod infestations and reduction of vector capacity to transmit pathogens that impact human and animal health. Recently, Cupp et al. [33] demonstrated that horn flies fed on cattle immunized with the anti-clotting factor thrombostasin, took smaller blood meals and the egg development was delayed. Although other molecules have been proposed as vaccine candidates against horn flies [16,34,35], further research is needed to identify new vaccine candidates for effective control of horn fly infestations.

Recently, RNA interference (RNAi) was proposed as a method to identify candidate tick protective antigens [36] and was used for the screening of tick genes with potential applications in vaccine development [37-39].

The aim of this study was to conduct a functional genomics study in female horn flies using Expressed Sequence Tags (EST) analysis and RNAi. The results of this study will advance the molecular characterization of this important ectoparasite and suggested candidate protective antigens for the development of vaccines for the control of horn fly infestations.

## Results

### Assembly and annotation of female horn fly Expressed Sequence Tags (ESTs)

A cDNA library was made from whole abdominal tissues collected from partially fed adult female horn flies. From 2,462 sequenced ESTs, 302 and 2,160 were low

and high quality ESTs, respectively (Table 1). Empty or vector ESTs were not obtained.


Since the female horn fly cDNA library was not normalized, the EST distribution per contig was quantified to determine the redundancy level of our EST dataset. High quality ESTs were assembled into 992 unigenes (178 contigs and 814 singlets) (Table 1; Additional file 1: Table S1), representing 46% novelty (unigenes/assembled ESTs) in our dataset. ESTs (814) present as singleton sequences represented 82% of all unigenes, while 72 unigenes (7%) contained only two ESTs. On average, the number of ESTs per unigene was 2.2, which suggested a low diversity in our dataset.

BLAST searches to TrEMBL and Swiss-Prot databases assigned 367 proteins to molecular function Gene Ontology (GO) terms (Figure 1). One hundred unigenes (10%) containing 535 ESTs (25%) corresponded to serine proteases. Other molecular functions represented in the unigenes included those involved in cell metabolism, mitochondrial function, transcription and translation, transport, chromatin structure, vitellogenesis, cytoskeleton, DNA replication, cell response to stress and infection, cell proliferation and cell-cell interactions, intracellular trafficking and secretion, and development (Figure 1). Of the 367 unigenes with molecular function GO assignments, 184 could be assigned to Clusters of Orthologous Groups of proteins (COG) (Figure 2). The COG comprising posttranslational modification, protein turnover and chaperones contained 40% of proteins with COG assignments, followed by translation, ribosomal structure and biogenesis (17%) and energy production and conversion (12%) (Figure 2).

A relatively large set of 449 unigenes (45%) lacked any significant sequence similarity (Blast  $E$  values  $> 10^{-5}$ ) to any sequence available in the public databases. Of all the 543 unigenes with significant sequence similarity to


**Table 1 Statistics of horn fly EST assembly**

Number of sequences	2,462
Mean length $\pm$ S.D. before vector stripping	877 $\pm$ 44 bp
Mean length $\pm$ S.D. after vector stripping	653 $\pm$ 32 bp
High quality EST reads	2,160
Assembled ESTs	2,160
No. unigenes	992
Mean length $\pm$ S.D.	758 $\pm$ 46 bp
No. unigenes with more than 20 ESTs	13
No. unigenes with 5-20 ESTs	44
No. unigenes with less than 5 ESTs	935
No. contigs	178
No. singlets	814
Novelty (unigenes/assembled ESTs)	46%
Redundancy (1-Novelsy)	54%



previously published sequences, 88.8% were most similar to Diptera (*Drosophila* spp., *H. irritans*, *Musca domestica*, *Lucilia cuprina*, *Glossina* spp., *Stomoxys calcitrans*, *Aedes aegypti*, *Sarcophaga* spp., *Phlebotomus papatasi*, *Chrysomya bezziana*, *Anopheles gambiae*, *Ceratitis stricta*, *Trichopalpus fraterna*, *Automola atomaria*, *Nanna tibiella*, *Bactrocera dorsalis*, *Lutzomyia longipalpis*, *Eristalinus punctulatus*, and *Ophiomyia* sp.), 1.5% to other insect species (*Spodoptera frugiperda*, *Tribolium castaneum*, *Gryllus bimaculatus*, *Lononia obliqua*, *Nasonia vitripennis*, and *Lymantria dispar*), 5.5% to other eukaryotic organisms, and 4.2% to microorganisms (Table 2).

Thirteen unigenes assembled from 505 sequence reads, contained more than 20 ESTs, most probably representing transcripts with highest abundance in abdominal tissues of partially fed female horn flies (Table 3). As expected from the results of the annotation of the entire EST dataset, 10 (77%) of these unigenes corresponded to serine proteases (Table 3). The second largest group of ESTs was derived from mitochondrial transcripts (Table 3). The analysis of serine protease unigene sequences showed that although some of them may be paralogs (for example unigenes 1-2 and 3-5; Table 3), other probably reflect sequence polymorphisms within the horn fly population because they



had 97%-98% nucleotide sequence identity (for example between unigenes 3-5; Table 3).

### Functional characterization of horn fly ESTs by RNAi

For functional genomics studies, selected unigene functional groups were used in RNAi experiments in female horn flies (Table 4). These groups included serine protease, protease inhibitor, vitellogenin (VTG), ubiquitination, ferritin (FER), vacuolar (H<sup>+</sup>)-ATPase (vATPase), proteasome component, immune response and 5'-nucleotidase (5'-NUC) ESTs and were selected based on their putative function in insect biology and previous results of RNAi experiments in other arthropods (see Discussion). As controls, ESTs with sequence identity to Nora virus and *Wolbachia* endosymbionts were selected (Table 4). The injection of these control dsRNAs did not affect horn fly mortality ( $\beta = -0.01$ , Wald  $\chi^2 = 0.01$ ,  $P = 0.91$ ) and oviposition ( $P > 0.05$ ) when compared to buffer-injected flies in 14 independent RNAi experiments (Table 5), thus supporting their use as controls. Significant gene knockdown was obtained for at least one targeted unigene sequence on each group except for the serine protease group 1 in which significant gene expression silencing was not obtained for any of the unigenes included in the analysis (Table 5). For some sequences, gene knockdown was observed as early as 6 h post-injection (hpi) and lasted at least until 36 hpi (Table 5). For other sequences in groups 8 and 9,

gene knockdown was not detected until after 12 hpi (Table 5). In most cases, gene expression silencing was higher than 70% when compared to the control group (Table 5).

To analyze RNAi off-target effects, the expression of genes not targeted by the injected dsRNA was analyzed at 12 hpi in functional groups 7-9 (Figure 3). The results showed that the expression of genes not targeted by the injected dsRNA was silenced in all three groups analyzed (Figure 3), thus suggesting RNAi off-target effects in horn flies. Pairwise sequence alignments identified regions with homology  $\geq 11$  bp in some sequences (Figure 4). However, only one region had 21 bp homology between unigene sequences 13\_D07 and 7\_A04 (Figure 4).

Injection of dsRNAs in the serine protease and ubiquitination functional groups did not affect fly mortality ( $\beta = 0.09$ , Wald  $\chi^2 = 2.68$ ,  $P = 0.10$  and  $\beta = 0.07$ , Wald  $\chi^2 = 3.60$ ,  $P = 0.08$ , respectively) or oviposition ( $P > 0.05$ ) when compared to controls (Table 5). The knockdown of a protease inhibitor gene (elastase) resulted in higher fly mortality ( $\beta = 0.60$ , Wald  $\chi^2 = 13.35$ ,  $P = 0.0002$ ) but did not affect oviposition ( $P > 0.05$ ) when compared to controls (Table 5). VTG-2 and proteasome component genes knockdown did not affect fly mortality ( $\beta = 0.03$ , Wald  $\chi^2 = 1.42$ ,  $P = 0.22$  and  $\beta = 0.13$ , Wald  $\chi^2 = 2.59$ ,  $P = 0.107$ , respectively) but significantly ( $P < 0.005$ ) reduced oviposition (Table 5). When the expression of immune response and 5'-NUC genes was silenced,

**Table 2 Distribution of annotated unigenes to different species**

Species	Total No. of unigenes	Ratio (%)
All organisms	543	100
Insects (Diptera)	482	88.8
<i>Drosophila</i> spp.	289	53.2
<i>H. irritans</i>	89	16.4
<i>Musca domestica</i>	22	4.0
<i>Lucilia cuprina</i>	16	2.9
<i>Glossina</i> spp.	15	2.8
<i>Stomoxys calcitrans</i>	14	2.6
<i>Aedes aegypti</i>	11	2.0
<i>Sarcophaga</i> spp.	9	1.7
<i>Phlebotomus papatasi</i>	3	0.6
<i>Chrysomya bezziana</i>	3	0.6
<i>Anopheles gambiae</i>	3	0.6
<i>Ceratitis stricta</i>	1	0.2
<i>Trichopalpus fraterna</i>	1	0.2
<i>Automola atomaria</i>	1	0.2
<i>Nanna tibiella</i>	1	0.2
<i>Bactrocera dorsalis</i>	1	0.2
<i>Lutzomyia longipalpis</i>	1	0.2
<i>Eristalinus punctulatus</i>	1	0.2
<i>Ophiomyia</i> sp.	1	0.2
Insects (Other)	8	1.5
<i>Tribolium castaneum</i>	2	0.4
<i>Gryllus bimaculatus</i>	2	0.4
<i>Spodoptera frugiperda</i>	1	0.2
<i>Lononia obliqua</i>	1	0.2
<i>Nasonia vitripennis</i>	1	0.2
<i>Lymantria dispar</i>	1	0.2
Other eukaryotes	30	5.5
Microorganisms	23	4.2

Over 90% of best hits matched to insects. Only less than 10% of all best hits matched to other eukaryotes and microorganisms.

higher fly mortality ( $\beta = -0.46$ , Wald  $\chi^2 = 7.39$ ,  $P = 0.006$  and  $\beta = 0.35$ , Wald  $\chi^2 = 4.65$ ,  $P = 0.03$ , respectively) and reduced oviposition ( $P < 0.005$ ) were obtained when compared to controls (Table 5). Interestingly, knockdown of FER light chain and vATPase genes resulted in lower fly mortality ( $\beta = 0.21$ , Wald  $\chi^2 = 5.12$ ,  $P = 0.02$  and  $\beta = -0.16$ , Wald  $\chi^2 = 14.70$ ,  $P = 0.0001$ , respectively), and 6 and 16-fold decrease in oviposition ( $P < 0.005$ ) when compared to control dsRNA-injected flies (Table 5).

## Discussion

The effective control of horn fly infestations requires the design of new control strategies. Genomics and functional genomics studies are important to understand basic biological questions and to identify new targets for

improved control strategies. Recently, gene expression analysis was reported in horn fly embryos, larvae and adult females [19,35]. However, this is the first report of functional genomics studies in this species.

ESTs sequenced and assembled in this study provided new sequence information for horn fly. The assembled unigenes without sequence similarity to sequences in public databases probably represented unique transcripts for horn fly or corresponded to proteins that have not yet been identified in related organisms due to incomplete genomic information. However, it cannot be excluded that the identified ESTs represent parts of known proteins whose similarities are located in parts of the sequence that are not covered by the analyzed ESTs.

The number of ESTs assembled into a certain unigene roughly reflected the relative abundance of corresponding mRNAs since the cDNA library from female horn flies used in this study was not normalized. We found that 100 unigenes, containing 25% of the ESTs, corresponded to serine proteases, indicating that this group represented the most abundantly expressed genes in abdominal tissues of partially fed female horn flies. In fact, the unigenes with the largest number of ESTs represented members of the serine protease family, thus suggesting that posttranslational modification and protein turnover were highly active in partially fed female flies.

The high proportion of ESTs present as singleton sequences when compared to contigs reflected a low diversity in our dataset, probably due to the presence of paralogs and sequence polymorphisms for some unigenes. In fact, sequence analysis of serine protease unigenes makes at this point difficult to discriminate between paralogs and ESTs representing sequence polymorphisms within the horn fly population.

RNAi was used to functionally characterize selected horn fly genes in adult female flies. To our knowledge, this is the first report of RNAi in horn flies. RNAi has been used to study gene function in insects and other arthropods [37,40-49] and to screen for candidate protective antigens in ticks [36-39]. Although with some fly mortality probably due to dsRNA injection with a Hamilton syringe, the RNAi method used here produced reproducible results in female horn flies. The failure to demonstrate gene knockdown for some sequences in serine protease and other functional groups studied could be due to unknown factors affecting RNAi in horn flies, because gene expression silencing did not occur until after 24-36 hpi for these genes or due to the existence of paralogs in these groups that affected the efficacy of gene knockdown. Gene silencing mediated by RNAi depends on short interfering RNAs (siRNAs) and micro RNAs (miRNAs). These RNAs have unique features, namely a defined size of 19-21 pb, and characteristic two-nucleotide single-stranded 3' overhangs and 5' monophosphate



**Table 3 Transcripts with highest abundance in abdominal tissues of partially fed female horn flies**

Unigene No. [GenBank accession number]	Number of clustered ESTs	Unigene annotation		
		Sequence identity	Clusters of Orthologous Groups of proteins (COG)	Molecular function Gene Ontology (GO) term
1 [HO004732]	21	<i>L. cuprina</i> clone sbasp9 serine proteinase	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
2 [HO004733]	21	<i>L. cuprina</i> clone sbasp9 serine proteinase	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
3 [HO004734]	22	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
4 [HO004735]	25	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
5 [HO004736]	25	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
6 [HO004737]	26	<i>D. pseudoobscura</i> GA21163-PA (Dpse\GA21163)	Amino acid transport and metabolism	Zinc carboxypeptidase
7 [HO004738]	28	<i>H. irritans</i> serine protease mRNA	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
8 [HO004739]	30	<i>L. cuprina</i> clone sbasp9 serine proteinase	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
9 [HO004740]	33	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
10 [HO004741]	37	<i>H. irritans</i> mitochondrion, complete genome		Mitochondria
11 [HO004742]	40	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
12 [HO004743]	90	<i>H. irritans</i> serine protease	Posttranslational modification, protein turnover, chaperones	Secreted trypsin-like serine protease
13 [HO004744]	107	<i>H. irritans</i> mitochondrion, complete genome		Mitochondria

groups [50]. Although RNAi off-target effects were shown in horn flies, most sequence alignments resulted in homology regions of 11 bp only and in some cases no homology  $\geq 11$  bp was found. These results suggested differences in RNAi specificity and sensitivity, a fact that needs to be fully characterized to understand and

efficiently use RNAi in horn flies and other organisms [37,48,51].

The aim of this study was to conduct a functional genomics study in female horn flies combining EST analysis with RNAi. Therefore, we will focus the discussion on unigene functional groups characterized by RNAi.

**Table 4 Horn fly unigene functional groups selected for RNAi**

Group N°	Functional group	N° of unigenes <sup>a</sup>	Unigenes selected for RNAi <sup>b</sup>
1	Serine protease	100	5, 10, 14, 19, 42, 90, 224, 230
2	Protease inhibitor	2	2_B12, 24_H02
3	Vitellogenin	13	7, 20, 37, 76, 89, 145, 176, 7_D07
4	Ubiquitination	5	84, 146, 4_E04, 5_G03, 7_B08
5	Ferritin	5	26, 39, 154, 156, 10_A09
6	vATPase	3	7_F08, 9_A08, 17_H03
7	Proteasome component	3	6_G04, 7_A04, 12_H09
8	Immune response	2	6_F11, 10_G05
9	5'-nucleotidase	1	13_D07
10 (negative control)	Infectious agents:	3	191
	Nora virus	3	2_E12
	<i>Wolbachia</i> endosymbionts		

<sup>a</sup>Number of unigenes grouped into this category.

<sup>b</sup>When the number of unigenes in the functional group was greater than 5, unigenes were selected for RNAi to include all different genes present in this category. Unigenes GenBank accession numbers are shown in Table 6.

**Table 5 Results of RNAi experiments in female horn flies**

Group No. <sup>a</sup>	Cumulative percent mortality			Oviposition (eggs per survived fly) <sup>b</sup>	Expression silencing (average% ± SD) with respect to group 10 control <sup>c</sup>				dsRNA injected <sup>d</sup>
	12 hpi	24 hpi	36 hpi		6 hpi	12 hpi	24 hpi	36 hpi	
1 (Serine protease)	57 ± 6	69 ± 11	78 ± 2	0.88 ± 0.23	ND	ND	57 ± 50 0 ± 0	ND	90 230
2* (Protease inhibitor)	73 ± 11	84 ± 7	94 ± 1	2.56 ± 1.80	ND	ND	83 ± 3**	ND	2_B12
3 (Vitellogenin)	55 ± 11	67 ± 8	74 ± 8	0.31 ± 0.9**	ND	ND	0 ± 0 0 ± 0 100 ± 2*	ND	7 20 76
4 (Ubiquitination)	42 ± 12	67 ± 11	83 ± 16	0.64 ± 0.63	ND	ND	92 ± 6* 70 ± 24 46 ± 43	ND	84 146 4_E04
5* (Ferritin)	24 ± 23	39 ± 37	47 ± 41	0.22 ± 0.17**	ND	ND	68 ± 13* 86 ± 86	ND	26 154
6* (vATPase)	17 ± 13	24 ± 11	34 ± 3	0.08 ± 0.05**	ND	ND	78 ± 6* 100 ± 1* 99 ± 8*	ND	7_F08 9_A08 17_H03
7 (Proteasome component)	64 ± 3	76 ± 8	84 ± 11	0.38 ± 0.03**	98 ± 2* 100 ± 6*	100 ± 2* 67 ± 9*	ND ND	0 ± 0 ND	7_A04 12_H09
8* (Immune response)	66 ± 7	ND	99 ± 8	0.23 ± 0.09**	100 ± 8* 0 ± 0	92 ± 5 0 ± 0	ND ND	96 ± 9* 98 ± 8*	6_F11 10_G05
9* (5'-nucleotidase)	50 ± 11	ND	91 ± 22	0.12 ± 0.09**	0 ± 0	98 ± 9*	ND	70 ± 7*	13_D07
10 (negative control)	45 ± 24	57 ± 30	64 ± 27	1.26 ± 0.90	—	—	—	—	191 2_E12
Injection buffer	46 ± 28	57 ± 30	65 ± 29	1.50 ± 1.01	0 ± 0	0 ± 0	0 ± 0	0 ± 0	None

<sup>a</sup>The average ± S.D. of two independent RNAi experiments for each of the test groups 1-9 and 14 experiments for the negative control and injection buffer groups is shown. One hundred flies were used on each RNAi experiment. Cumulative percent mortality was evaluated in female horn flies at 12, 24 and 36 hpi. Survival curves (temporal rates of mortality) were compared between different treatments and the control group 10 using Cox Proportional Hazards Survival Regression analysis (\*P < 0.05). Abbreviation: ND, not determined.

<sup>b</sup>Oviposition data from test groups 1-9 and the injection buffer control were compared with the unrelated dsRNA-injected control group 10 by Student's t-test (\*\*P < 0.005).

<sup>c</sup>For analysis of expression silencing, the mRNA levels of each knockdown gene were determined by real-time RT-PCR in 4 individual flies each at 6, 12 and 24 or 36 hpi. Percent expression silencing shown at different hpi corresponds to each individual gene targeted by dsRNA injection. The mRNA levels were normalized against horn fly 16S rRNA and the mean of the duplicate normalized Ct values from test dsRNA-injected groups 1-9 and in the injection buffer control were compared with the unrelated dsRNA-injected control group 10 by Student's t-test (\*P < 0.05, \*\*P < 0.005).


<sup>d</sup>Unigenes GenBank accession numbers are shown in Table 6.

## Serine proteases

Serine proteases are a group of endopeptidases involved in several processes such as digestion, immune response, blood clotting and inflammation. In female horn flies, 10% of the assembled unigenes, containing more than 500 ESTs, were identified as serine proteases. In agreement with these results, Guerrero et al. [19] recently showed that serine proteases are differentially expressed in fly adult stages when compared to larvae. Significant gene knockdown was not obtained for any of the genes targeted by dsRNA injection in this group. Consequently, RNAi did not affect fly mortality or oviposition. In other arthropods, silencing of serine proteases expression by RNAi showed that these proteins are involved in blood digestion, oocyte maturation, development and immune response [40,42-44,46,52-55].


## Protease inhibitors

The protease inhibitor genes identified in female horn flies corresponded to serpins, inhibitors of serine proteases and thus involved in the same biological processes discussed before for serine proteases. A horn fly serine protease inhibitor gene was previously cloned and characterized, suggesting that these genes may be involved in the control of fly endogenous and pathogen proteases [56,57]. In mosquitoes, serpin RNAi affected insect immune response [58]. The elastase inhibitor gene knockdown significantly increased horn fly mortality at 12, 24 and 36 hpi. Thus, the effect of elastase inhibitor RNAi described here in horn flies may be the result of impaired fly protease control and/or the effect of increased susceptibility to persistent pathogen infections resulting from diminished immune response.



Vitellogenin

VTGs constitute a multigene superfamily encoding for egg yolk precursor proteins expressed in the females of arthropods and other oviparous organisms [59]. In cockroaches, ants and ticks, knockdown of VTG receptor (VTG-R), essential for VTG uptake into developing oocytes, disrupts egg formation [41,60-62]. In honeybees, silencing of VTG expression by RNAi affects honeybee workers developmental behavior [63,64]. Similar to results of VTG-R knockdown in other arthropods, silencing of VTG-2 expression in horn flies reduced oviposition in 4-fold when compared to controls.



Ubiquitination

Ubiquitination is a post-translational modification carried out by a set of enzymes that affect protein proteasomal degradation, stability, function, and intracellular localization [65]. In this functional group, horn fly genes involved in the ubiquitination pathway such as ubiquitin-1 (UBQ-1), UBQ-protein ligase, and UBQ hydrolase were included. In this group, only the UBQ-protein ligase expression was significantly silenced after RNAi. Although UBQ-protein ligase has been shown to regulate apoptosis in *Drosophila* [66], knockdown of this gene did not affect horn fly mortality or oviposition. Thus, it may be possible that the phenotype resulting from silencing the UBQ-protein ligase expression was not evident in horn flies under our experimental conditions. Additionally, knockdown of other ubiquitination genes may be required in order to have a significant effect on horn fly mortality and oviposition. For example, UBQ knockdown in ticks causes mortality and reduces oviposition and *Anaplasma marginale* infection/multiplication in the guts [39,67].

Ferritin

FER is the main protein for intracellular iron storage and consists of 2 types of subunits, a heavy (ferroxidase sites) and a light chain (nucleation sites) [68]. FER light



(3 unigenes containing 5 ESTs) and heavy (2 unigenes containing 4 ESTs) chains were not among the most abundant ESTs identified in female horn flies. However, Guerrero et al. [35] found FER light chain as one of the most abundant transcripts in horn fly larvae. These results suggested differences in the FER expression between horn fly larvae and adult females. FER light chain knockdown in horn flies significantly reduced oviposition (6-fold with respect to controls), but surprisingly fly mortality was reduced when compared to controls. In ticks, FER RNAi reduces not only oviposition but also feeding and *A. marginale* infection levels in IDE8 cells [67,68].

#### vATPase

vATPase is a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles and has been shown to be required for the normal function of the Golgi complex, endoplasmic reticulum, vacuoles and endocytotic and exocytotic vesicles [69]. vATPase was also implicated in immunity [70]. Guerrero et al. [35] identified vATPase as one of the most abundant transcripts in horn fly larvae. However, in adult females, only 3 ATPase unigenes were assembled with one EST each, those suggesting like previously for FER, differences in the vATPase expression between horn fly larvae and adult females. Genetic knockout of vATPase subunits resulted in lethal phenotypes in fruit flies (*Drosophila melanogaster*), flour beetles (*T. castaneum*), pea aphids (*Acyrtosiphon pisum*), and tobacco hornworms (*Manduca sexta*) [69,71] and reduced influenza virus replication in *Drosophila* cells [72]. RNAi of vATPase expression in ticks resulted in testis and salivary gland degeneration, suggesting a role for this molecule in the function of these organs [73] and reduced *A. marginale* infection in *Dermacentor variabilis* tick guts but not pathogen multiplication in IDE8 tick cells [67]. vATPase knockdown in horn flies resulted in 16-fold reduction in oviposition but, as with FER light chain, fly mortality was reduced when compared to controls. These results suggested that despite the important function of vATPase in all arthropods, developmental stage-specific and species-specific differences might exist that could explain the results obtained after gene knockdown in horn flies.

#### Proteasome component

Proteasomes are large protein complexes involved in protein proteolysis that are functionally related to ubiquitination and thus essential for eukaryotic cells [74]. Experiments in *D. melanogaster* showed that knockdown of proteasome subunits leads to increased levels of ubiquitin conjugates, cell cycle defects, DNA overreplication, and apoptosis [74,75]. In tick cells, 26S proteasome

knockdown resulted in lower *A. marginale* infection levels when compared to controls but did not affect tick survival, feeding and reproduction [67]. However, based on the essential proteasome function in eukaryotic cells, it was not surprising to observe a decrease in oviposition in horn flies injected with proteasome components dsRNAs targeting proteasome subunit beta (two unigenes) and proteasome maturation protein (one unigene). As previously shown in *D. melanogaster* [74,75], proteasome subunits knockdown in horn flies may affect cell cycle and DNA replication thus resulting in reduced oviposition.

#### Immune response

Innate immune response is essential for insect survival. Only two unigenes were assembled into this category and knockdown in female horn flies. Assembled unigenes encoded for putative T-cell immunomodulatory protein and RNase L inhibitor. Silencing of these genes resulted in higher horn fly mortality and lower oviposition when compared to controls. These RNAi results may be due to an effect of gene knockdown on increased susceptibility to persistent pathogen infections resulting from impaired immune response in horn flies. Knockdown of immune response genes may affect the mechanisms involved in the control of persistent infections such as those caused by Nora virus and *Wolbachia* spp. [76-78], which could affect horn fly mortality and oviposition. RNAi knockdown of immune response genes in other arthropods results in increased mortality and higher pathogen infection levels [79-81].

#### 5'-nucleotidase

5'-NUC and other ectonucleotidases control the levels of extracellular nucleotides and nucleosides that act as signaling molecules involved in a wide spectrum of biological effects [82]. 5'-NUC is commonly expressed in the salivary glands of blood-sucking ectoparasites [83-88]. Herein, as previously shown in ticks [36], 5'-NUC knockdown resulted in higher fly mortality and lower oviposition when compared to controls. As in other organisms, these results suggested an essential function for 5'-NUC in horn fly females.

#### Conclusions

In summary, a cDNA library was constructed from whole abdominal tissues collected from partially fed adult female horn flies and 2,160 high quality ESTs were sequenced and assembled into 992 unigenes (178 contigs and 814 singlets) representing molecular functions such as serine proteases, cell metabolism, mitochondrial function, transcription and translation, transport, chromatin structure, vitellogenesis, cytoskeleton, DNA replication, cell response to stress and infection, cell proliferation and cell-cell interactions, intracellular

trafficking and secretion, and development. A method was developed for RNAi that produced reproducible results in horn flies. Functional analyses by RNAi showed the effect of some genes on horn fly mortality and oviposition. These results advanced the molecular characterization of this important ectoparasite and suggested candidate protective antigens for the development of vaccines for the control of horn fly infestations. Based on RNAi results, some of the candidate antigens to be considered for cattle vaccination experiments against horn flies include those within VTG, immune response and 5'-NUC functional groups.

## Methods

### Rearing of horn flies

*H. irritans* were reared under laboratory conditions as reported by Schmidt et al. [89]. A horn fly colony was established with flies originally collected in a cattle farm close to Ciudad Victoria, Tamaulipas, Mexico. About 2,000 flies were collected from 2 infested animals and transported in a 20 × 30 cm mosquito netting aluminum cage. Flies were allowed to lay eggs over a water container during 12 h. Eggs were collected and incubated into fresh bovine feces during 5 days. Pupae were collected and placed in Petri dishes located inside mosquito netting aluminum cages for molting into adult flies. After molting, flies were fed twice a day using pieces of cotton impregnated with fresh defibrinated bovine blood obtained from a naive cow. All the horn fly developmental phases were kept under a photoperiod of 12 h light: 12 h darkness at 28-32°C and 70-80% relative humidity [89].

### Analysis of expressed sequence tags (ESTs) in adult female horn flies

Total RNA was isolated from whole abdominal tissues collected from 1,500 partially fed adult female horn flies using Trizol (Sigma, St. Louis, MO, EUA). The cDNA library was synthesized using the SMART™ cDNA Library Construction Kit (Clontech, Mountain View, CA, USA) at Creative Biolabs (Port Jefferson Station, NY, USA; <http://www.creativebiolabs.com>). cDNAs were cloned into the pBluescript II SK vector (Agilent Technologies, Inc., Santa Clara, CA, USA). The library had more than  $1 \times 10^6$  primary clones, with >90% recombinants with inserts >500 bp (average cDNA length >1,000 bp). A total of 2,462 ESTs were 5' sequenced (Creative Biolabs). The cDNA Annotation System software (CAS; Bioinformatics and Scientific IT Program (BSIP), Office of Technology Information Systems (OTIS), National Institute of Allergy and Infectious Diseases (NIAID), Bethesda, MD, USA) <http://exon.niaid.nih.gov> was used for automated sequence clean up, assembly, blasting against multiple sequence

databases (ncbi non-redundant nucleotide and protein sequence databases, *H. irritans* EST sequences [35] and databases of mosquito- and tick-specific sequences <http://www.ncbi.nlm.nih.gov/>; <http://www.vectorbase.org/index.php>) and Gene Ontology (GO) assignments. Comparison with the ncbi Clusters of Orthologous Groups of proteins (COG; <http://www.ncbi.nlm.nih.gov/COG>) was also performed. Nucleotide sequences were aligned using the program AlignX (Vector NTI Suite V 5.5, InforMax, North Bethesda, MD, USA). Gene sequences were deposited in the GenBank with accession numbers HO000420-HO001165 and HO004499-HO004744.

### RNAi in adult female horn flies

Oligonucleotide primers (pBLUET75: 5'-*TAATACGACT CACTATAGGGTACTTCGAGGTGACGGTATCGAT*-3' and pBLUET73: 5'-*TAATACGACTCACTATAGGGTACT-CAATTAACCTCACTAAAGGGA*-3') were synthesized specific for vector DNA sequences flanking the horn fly cDNA insert and containing T7 promoter sequences (in italics) for *in vitro* transcription and synthesis of dsRNA. PCR reactions were performed from individual or pooled cDNA clones (when more than one unigene was included in the functional group analyzed; Table 2) using the Access RT-PCR system (Promega, Madison, WI, USA) in a 50 µl reaction mixture. The resultant amplicons were purified using the Wizard 96-well PCR purification system (Promega). *In vitro* transcription and purification of dsRNA was done using the Megascript RNAi kit (Ambion, Austin, TX, USA). The dsRNA was quantified by spectrometry. Adult partially fed female flies were injected with approximately 0.1 µl of dsRNA ( $1 \times 10^9$ - $1 \times 10^{11}$  molecules per µl) in the abdominal segment. The injections were done with a Hamilton syringe with a 1 inch, 33 gauge needle. Control flies were injected with unrelated dsRNA or injection buffer (10 mM Tris-HCl, pH 7, 1 mM EDTA) (Table 5). One hundred flies were used in each group. After injection with dsRNA, female flies were kept in petri dishes for one hour and then transferred to wired 20 × 30 cm boxes. Flies were fed using impregnated cotton with fresh defibrinated blood obtained from a naive cow and reared as described before. Fly mortality was evaluated at 12, 24 and 36 hpi. Survival curves (temporal rates of mortality) were compared between different treatments and controls using Cox Proportional Hazards Survival Regression analysis (SPSS Inc., Chicago, IL, USA). Oviposition (number of eggs per survived fly) was also evaluated and the results in test dsRNA-injected groups and in the injection buffer control were compared with the unrelated dsRNA-injected control group by Student's t-test ( $P = 0.05$ ).

**Table 6 Primer sets and real-time RT-PCR conditions used for analysis of selected *H. irritans* ESTs**

Unigene ID (Genbank accession number)	Gene description	Upstream/downstream primer sequences (5'-3')	PCR annealing conditions
13_D07 (HO000820)	5'-nucleotidase	AGTGGACAAATGTCCCGAAG AGCATTGGGGTTTGAATGAG	55°C, 30 s
6_F11 (HO000609)	T-cell immunomodulatory protein	CCGGTGACTTTGATGGAGAT GATAATGGCTCCCTTTTGGT	55°C, 30 s
10_G05 (HO000738)	RNase L inhibitor	GCCGATCGTGTTATTGTCTT CCGGATCGTTTTTGTTCAGT	55°C, 30 s
7_A04 (HO000619)	Proteasome subunit beta	CAGGCGAGGTCCATTATTGT AGTGCGCGACCTCAAGTAGT	55°C, 30 s
12_H09 (HO000808)	Proteasome maturation protein	GAGGAATCGTGAGGGTTTGA ACATGGGGTTGTCTCGATAAA	55°C, 30 s
7_F08 (HO000644)	vATPase subunit d	TGTTTTCCGTCAACAGTCA GGCACAACCTCCAAGTAA	60°C, 30 s
9_A08 (HO000679)	vATPase subunit f	TGTTGGATTCTTGCTTGGTG GGCACTGGTGATGTATGTGC	60°C, 30 s
17_H03 (HO001053)	vATPase proteolipid subunit	GTCCAGCCAGACTGTGATGA AATCAATCGCGGACAAAAAC	60°C, 30 s
26 (HO004524)	Ferritin light chain	TGATCATGTTGAACCCGAGA CGGCTGGTCAATTTCTTGAT	60°C, 30 s
154 (HO004652)	Ferritin heavy chain	GTTGTTGCCCTGCTGTATT TGAAAAGTGGGCTCCCATAG	60°C, 30 s
84 (HO004582)	Ubiquitin-protein ligase	TCGCATCTGTTTGGATGTGT CGGGAAAACCTTTGAGTCCA	60°C, 30 s
146 (HO004644)	Ubiquitin family (UBQ-1)	CCCGACCAACAACGTTTAAT CGACGAAGACGGTGAATTTT	60°C, 30 s
4_E04 (HO000538)	Ubiquitin carboxyl-terminal hydrolase	AGCCAGAGATGTTGGAATGG TCGATGTAAATTGCCGCATA	60°C, 30 s
7 (HO004505)	Vitellogenin 3	GAGCTTTTTGCGTTGTAGCC ACAAAAGTGGGAGCAACACC	60°C, 30 s
20 (HO004518)	Vitellogenin 1	GAGCTTTTTGCGTTGTAGCC ACAAAAGTGGGAGCAACACC	60°C, 30 s
76 (HO004574)	Vitellogenin 2	ACGGCCGGTTGTGAGATTAT AGCATCTTTTCGGTCTTGC	60°C, 30 s
2_B12 (HO000479)	Elastase inhibitor	CAAGGGTGAATGGGAAAAGA TAAAGGCCTTCACGTTCTTG	60°C, 30 s
90 (HO004588)	Serine protease, midgut specific trypsin, secreted	TGCGTTATATCCGTTGGTG CTTTGTCAACGGCATAAGCA	60°C, 30 s
230 (HO004728)	Serine protease	TGGCTACAATGAATGCAAGC GGTTAGCACCAGGGAACGTA	60°C, 30 s
(FJ025436)	<i>H. irritans</i> 16S rRNA	TTTAAATGGCCGAGTATCC GATTATAGGGTCTTCTCGTCTTTT	60°C, 30 s

Gene expression silencing was evaluated in 4 individual flies each at 6, 12 and 36 or 24 hpi. The mRNA levels of each knockdown gene were determined using sequence-specific oligonucleotide primers (Table 6) and the iScript One-Step RT-PCR Kit with SYBR Green and the iQ5 thermal cycler (Bio-Rad, Hercules, CA, USA) following manufacturer's recommendations. A dissociation curve was run at the end of the reaction to ensure that only one amplicon was formed and that the amplicon denatured consistently in the same temperature range for every sample [90]. The mRNA levels were normalized against horn fly 16S rRNA (Table 4) using

the genNorm method (ddCT method as implemented by Bio-Rad iQ5 Standard Edition, Version 2.0) [91]. In all cases, the mean of the duplicate values was used and normalized Ct values from test dsRNA-injected groups and in the injection buffer control were compared with the unrelated dsRNA-injected control group by Student's t-test ( $P = 0.05$ ).

## Additional material

**Additional file 1: Table S1.** Assembling and analysis of *H. irritans* high quality ESTs.

# Acknowledgements

Joaquín Vicente (IREC, Spain) is acknowledged for conducting statistical analysis of fly survival curves. This research was supported by SAGARPA-CONACYT (project 12260) and SEP-CONACYT (project 25772), Mexico to CA, the INIA (project FAU2008-00014-00-00) and the University of Castilla La Mancha (project 00030072 "Ayudas para acciones de cooperación al desarrollo"), Spain to JF. Lorena Torres was supported by CONACYT, grant 42883 and Promep-UAT, Mexico. R.C. Galindo was funded by MEC, Spain. Publication of this paper was possible through the grant 151779, FOMIX, Tamaulipas, Mexico.

# Author details

<sup>1</sup>Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Tamaulipas, Km. 5 carretera Victoria-Mante, CP 87000 Ciudad Victoria, Tamaulipas, Mexico. <sup>2</sup>Instituto de Investigación en Recursos Cinegéticos IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13005 Ciudad Real, Spain. <sup>3</sup>Centro Nacional de Investigación Disciplinaria en Parasitología Veterinaria, Carretera Federal Cuernavaca-Cuautla 8534, Col Progreso, Jiutepec, Morelos. CP 62550, Mexico. <sup>4</sup>Departamento de Parasitología, Facultad de Medicina Veterinaria y Zootecnia, Universidad Nacional Autónoma de México, Ciudad Universitaria, Coyoacán, D.F. CP 04510, Mexico. <sup>5</sup>Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK 74078, USA.

# Authors' contributions

LT reared horn flies, prepared RNA samples, did the RNAi experiments and helped with drafting the manuscript. CA participated in coordination of the study, RNAi experiments and drafting of the manuscript. NA, RCG and CA conducted the real-time RT-PCR analyses. RCG helped with drafting the manuscript. JF participated in design and coordination of the study, analyzed EST and RNAi data and drafted the manuscript. RRC and HQR participated in coordination of the study. All authors read and approved the final manuscript.

Received: 7 June 2010 Accepted: 10 February 2011

Published: 10 February 2011

# References

1. Harwood RF, James MT: *Entomología Médica y Veterinaria. Uteha, Mexico*, DF 1993.
2. Barros AT, Guglielmon AA, Martins JR: *Mosca de los cuernos: Control sustentable y resistencia a los insecticidas. Red Ectopar* 2002, 1-10.
3. Quiroz RH: *Parasitología y Enfermedades Parasitarias de los Animales Domésticos*. Limusa. México, DF, 3 2005.
4. Cupp EW, Cupp MS, Ribeiro JM, Kunz SE: *Blood-feeding strategy of Haematobia irritans (Diptera: Muscidae)*. *J Med Entomol* 1998, 35:591-595.
5. Steelman CD, Brown MA, Gbur EE, Tolley G: *The effects of hair density of beef cattle on Haematobia irritans horn fly populations. Med Vet Entomol* 1997, 11:257-264.
6. Johnson NN, Mayer DG: *Estimation of the effects of buffalo fly (Haematobia irritans exigua) on the estimation of dairy cattle based on a meta-analysis of literature data. Med Vet Entomol* 1999, 13:372-376.
7. Kunz SE, Kemp DH: *Insecticides and acaricides: resistance and environmental impact. Rev Sci Tech* 1994, 13:1249-1286.
8. Cupp EW, Cupp MS, Ribeiro JM, Kunz SE: *Blood-feeding strategy of Haematobia irritans (Diptera: Muscidae)*. *J Med Entomol* 1998, 35:591-595.
9. Guglielmon AA, Gimeno E, Idriat J, Fisher WF, Volpogni MM, Quaino O, Anziani OS, Flores SG, Warnke O: *Skin lesions and cattle hide damage from Haematobia irritans infestations. Med Vet Entomol* 1999, 13:324-329.
10. Sinshaw A, Abebe G, Desquesnes M, Yoni W: *Biting flies and Tripanosoma vivax infection in three highland districts bordering lake Tana, Ethiopia. Vet Parasitol* 2006, 142:35-46.
11. Abril C, Nimmervoll H, Pilo P, Brodard I, Korczak B, Markus S: *Rapid diagnosis and quantification of Francisella tularensis in organs of naturally infected common squirrel monkeys (Saimiri sciureus)*. *Vet Microbiol* 2007, 127:203-208.
12. Gillespie BE, Owens WE, Nickerson SC, Oliver SP: *Deoxyribonucleic acid fingerprinting of Staphylococcus aureus from heifer mammary secretions from horn flies. J Dairy Sci* 1999, 82:1581-1585.
13. Spier SJ, Leutenegger CM, Carroll SP, Loye JE, Pusteria JB, Carpenter TE, Mihalji JE, Madigan JE: *Use of real-time polymerase chain reaction-based fluorogenic 5' nuclease assay to evaluate insect vectors of Corynebacterium pseudotuberculosis infections in horses. Amer J Vet Res* 2004, 65:829-834.
14. Rodríguez SD, García Ortiz MA, Jiménez Ocampo R, Vega y Murguía C: *Molecular epidemiology of bovine anaplasmosis with a particular focus in Mexico. Infect Genet Evol* 2009, 9:1092-1101.
15. Kunz SE, Schmidt CD: *The pyrethroid resistance problem in the horn fly. J Agric Entomol* 1985, 2:358-363.
16. Oyarzún MP, Quiroz A, Birkett MA: *Insecticide resistance in the horn fly: alternative control strategies. Med Vet Entomol* 2008, 22:188-202.
17. Kunz SE, Kemp DH: *Insecticides and acaricides: Resistance and environmental impact. Rev Sci Tech* 1994, 13:1249-1286.
18. Mochi DA, Monteiro AC, Ribeiro Machado AC, Yoshida L: *Efficiency of entomopathogenic fungi in the control of eggs and larvae of the horn fly Haematobia irritans (Diptera: Muscidae)*. *Vet Parasitol* 2009, 167:62-66.
19. Guerrero FD, Dowd SE, Sun Y, Saldivar L, Wiley GB, Macmill SL, Najar F, Roe BA, Foil LD: *Microarray analysis of female- and larval-specific gene expression in the horn fly (Diptera: Muscidae)*. *J Med Entomol* 2009, 46:257-270.
20. Willadsen P, Riding GA, McKenna RV, Kemp DH, Tellam RL, Nielsen JN, Lahstein J, Cobon GS, Gough JM: *Immunological control of a parasitic arthropod: identification of a protective antigen from Boophilus microplus. J Immunol* 1989, 143:1346-1351.
21. de la Fuente J, Almazán C, Canales M, Pérez de la Lastra JM, Kocan KM, Willadsen P: *A ten-year review of commercial vaccine performance for control of tick infestations on cattle. Anim Health Res Rev* 2007, 8:23-28.
22. de la Fuente J, Rodríguez M, Redondo M, Montero C, García-García JC, Méndez L, Serrano E, Valdés M, Enríquez A, Canales M, Ramos E, de Armas CA, Rey S, Rodríguez JL, Artiles M, García L: *Field studies and cost-effectiveness analysis of vaccination with Gavac™ against the cattle tick Boophilus microplus. Vaccine* 1998, 16:366-373.
23. de la Fuente J, Kocan KM: *Advances in the identification and characterization of protective antigens for development of recombinant vaccines against tick infestations. Expert Rev Vaccines* 2003, 2:583-593.
24. Valenzuela JG, Belkaid Y, Garfield MK, Mendez S, Kamhawi S, Rowton ED, Sacks DL, Ribeiro JM: *Toward a defined anti-Leishmania vaccine targeting vector antigens: characterization of a protective salivary protein. J Exp Med* 2001, 194:331-342.
25. Lal AA, Patterson PS, Sacchi JB, Vaughan JA, Paul C, Collins WE, Wirtz RA, Azad AF: *Anti-mosquito midgut antibodies block development of Plasmodium falciparum and Plasmodium vivax in multiple species of Anopheles mosquitoes and reduce vector fecundity and survivorship. Proc Natl Acad Sci USA* 2001, 98:5228-5233.
26. Almeida AP, Billingsley PF: *Induced immunity against the mosquito Anopheles stephensi (Diptera: Culicidae): effects of cell fraction antigens on survival, fecundity, and Plasmodium berghei (Eucoccidiidae: Plasmodiidae) transmission. J Med Entomol* 2002, 39:207-214.
27. Suneja A, Gulia M, Gakhar SK: *Blocking of malaria parasite development in mosquito and fecundity reduction by midgut antibodies in Anopheles stephensi (Diptera: Culicidae)*. *Arch Insect Biochem Physiol* 2003, 52:63-70.
28. Milleron RS, Ribeiro JM, Elnaime D, Soong L, Lanzaro G: *Negative effect of antibodies against maxadilan on the fitness of the sand fly vector of American visceral leishmaniasis. Am J Trop Med Hyg* 2004, 70:278-285.
29. Titus RG, Bishop JV, Mejia JS: *The immunomodulatory factors of arthropod saliva and the potential for these factors to serve as vaccine targets to prevent pathogen transmission. Parasite Immunol* 2006, 4:131-141.
30. Kedzierski L, Zhu Y, Handman E: *Leishmania vaccines: progress and problems. Parasitol* 2006, 133:S87-S112.
31. Saul A: *Mosquito stage, transmission blocking vaccines for malaria. Curr Opin Infect Dis* 2007, 5:476-481.
32. Canales M, Naranjo V, Almazán C, Molina R, Tsuruta SA, Szabó MPJ, Manzano-Roman R, Pérez de la Lastra JM, Kocan KM, Jiménez MI, Lucientes J, Villar M, de la Fuente J: *Conservation and immunogenicity of the mosquito ortholog of the tick protective antigen, subolesin. Parasitol Res* 2009, 105:97-111.
33. Cupp MS, Cupp EW, Navarre C, Wisniewski N, Brandt KS, Silver GM, Zhang D, Panangala V: *Evaluation of a recombinant salivary gland protein (thrombostasin) as a vaccine candidate to disrupt blood-feeding by horn flies. Vaccine* 2004, 22:2285-2297.
34. Wijffels G, Hughes S, Gough J, Allen J, Don A, Marshall K, Kay B, Kemp D: *Peritrophins of adult dipteran ectoparasites and their evaluation as vaccine antigens. Int J Parasitol* 1999, 29:1363-77.



35. Guerrero FD, Dowd SE, Nene VM, Foil LD: Expressed cDNAs from embryonic and larval stages of the horn fly (Diptera: Muscidae). *J Med Entomol* 2008, **45**:686-692.
36. de la Fuente J, Almazán C, Blouin EF, Naranjo V, Kocan KM: RNA interference screening in ticks for identification of protective antigens. *Parasitol Res* 2005, **96**:137-141.
37. de la Fuente J, Kocan KM, Almazán C, Blouin EF: RNA interference for the study and genetic manipulation of ticks. *Trends Parasitol* 2007, **23**:427-433.
38. de la Fuente J, Manzano-Roman R, Naranjo V, Kocan KM, Zivkovic Z, Blouin EF, Canales M, Almazán C, Galindo RC, Step DL, Villar M: Identification of protective antigens by RNA interference for control of the lone star tick, *Amblyomma americanum*. *Vaccine* 2010, **28**:1786-1795.
39. Almazán C, Lagunes R, Villar M, Canales M, Rosario-Cruz R, Jongejan F, de la Fuente J: Identification and characterization of *Rhipicephalus (Boophilus) microplus* candidate protective antigens for the control of cattle tick infestations. *Parasitol Res* 2010, **106**:471-479.
40. Miyoshi T, Tsuji N, Islam MK, Kamio T, Fujisaki K: Gene silencing of a cubilin-related serine proteinase from the hard tick *Haemaphysalis longicornis* by RNA interference. *J Vet Med Sci* 2004, **66**:1471-3.
41. Ciudad L, Piulachs MD, Bellés X: Systemic RNAi of the cockroach vitellogenin receptor results in a phenotype similar to that of the *Drosophila* yolkless mutant. *FEBS J* 2006, **273**:325-35, Erratum in: *FEBS J* 2006 Feb;273(3):671.
42. Kambris Z, Brun S, Jang IH, Nam HJ, Romeo Y, Takahashi K, Lee WJ, Ueda R, Lemaître B: *Drosophila* immunity: a large-scale in vivo RNAi screen identifies five serine proteases required for Toll activation. *Curr Biol* 2006, **16**:808-13.
43. Araujo RN, Campos IT, Tanaka AS, Santos A, Gontijo NF, Lehane MJ, Pereira MH: Brasiensin: A novel intestinal thrombin inhibitor from *Triatoma brasiliensis* (Hemiptera: Reduviidae) with an important role in blood intake. *Int J Parasitol* 2007, **37**:1351-8.
44. Wei Z, Yin Y, Zhang B, Wang Z, Peng G, Cao Y, Xia Y: Cloning of a novel protease required for the molting of *Locusta migratoria manilensis*. *Dev Growth Differ* 2007, **49**:611-21.
45. Chen XG, Mathur G, James AA: Gene expression studies in mosquitoes. *Adv Genet* 2008, **64**:19-50.
46. Isoe J, Rascón AA Jr, Kunz S, Miesfeld RL: Molecular genetic analysis of midgut serine proteases in *Aedes aegypti* mosquitoes. *Insect Biochem Mol Biol* 2009, **39**:903-12.
47. Krishnan P, Gireesh-Babu P, Rajendran KV, Chaudhari A: RNA interference-based therapeutics for shrimp viral diseases. *Dis Aquat Organ* 2009, **86**:263-72.
48. Bellés X: Beyond *Drosophila*: RNAi in vivo and functional genomics in insects. *Annu Rev Entomol* 2010, **55**:11-28.
49. Huvenne H, Smagghe G: Mechanisms of dsRNA uptake in insects and potential of RNAi for pest control: a review. *J Insect Physiol* 2010, **56**:227-35.
50. Lingel A, Sattler M: Novel modes of protein-RNA recognition in the RNAi pathway. *Curr Opin Struct Biol* 2005, **15**:107-15.
51. de la Fuente J, Maritz-Olivier C, Naranjo V, Ayoubi P, Nijhof AM, Almazán C, Canales M, Pérez de la Lastra JM, Galindo RC, Blouin EF, Gortazar C, Jongejan F, Kocan KM: Evidence of the role of tick subolesin in gene expression. *BMC Genomics* 2008, **9**:372.
52. Brackney DE, Foy BD, Olson KE: The effects of midgut serine proteases on dengue virus type 2 infectivity of *Aedes aegypti*. *Am J Trop Med Hyg* 2008, **79**:267-74.
53. Shah PK, Tripathi LP, Jensen LJ, Gahnm M, Mason C, Furlong EE, Rodrigues V, White KP, Bork P, Sowdhamini R: Enhanced function annotations for *Drosophila* serine proteases: a case study for systematic annotation of multi-member gene families. *Gene* 2008, **407**:199-215.
54. Marshall JL, Huestis DL, Hiromasa Y, Wheeler S, Oppert C, Marshall SA, Tomich JM, Oppert B: Identification, RNAi knockdown, and functional analysis of an ejaculate protein that mediates a postmating, prezygotic phenotype in a cricket. *PLoS One* 2009, **4**:e7537.
55. Amparyup P, Wiriyakaradecha K, Charoensapsri W, Tassanakajon A: A clip domain serine proteinase plays a role in antibacterial defense but is not required for prophenoloxidase activation in shrimp. *Dev Comp Immunol* 2010, **34**:168-76.
56. Azzolini SS, Santos JM, Souza AF, Torquato RJ, Hirata IY, Andreotti R, Tanaka AS: Purification, characterization, and cloning of a serine proteinase inhibitor from the ectoparasite *Haematobia irritans irritans* (Diptera: Muscidae). *Exp Parasitol* 2004, **106**:103-109.
57. Azzolini SS, Sasaki SD, Campos IT, S Torquato RJ, Juliano MA, Tanaka AS: The role of HiTI, a serine protease inhibitor from *Haematobia irritans irritans* (Diptera: Muscidae) in the control of fly and bacterial proteases. *Exp Parasitol* 2005, **111**:30-36.
58. Michel K, Suwanchaichinda C, Morlais I, Lambrechts L, Cohuet A, Awono-Ambene PH, Simard F, Fontenille D, Kanost MR, Kafatos FC: Increased melanizing activity in *Anopheles gambiae* does not affect development of *Plasmodium falciparum*. *Proc Natl Acad Sci USA* 2006, **103**:16858-63.
59. Byrne BM, Gruber M, Ab G: The evolution of egg yolk proteins. *Prog Biophys Mol Biol* 1989, **53**:33-69.
60. Mitchell RD, Ross E, Osgood C, Sonenshine DE, Donohue KV, Khalil SM, Thompson DM, Michael Roe R: Molecular characterization, tissue-specific expression and RNAi knockdown of the first vitellogenin receptor from a tick. *Insect Biochem Mol Biol* 2007, **37**:375-88.
61. Boldbaatar D, Battsetseg B, Matsuo T, Hatta T, Umeyama-Shirafuji R, Xuan X, Fujisaki K: Tick vitellogenin receptor reveals critical role in oocyte development and transovarial transmission of *Babesia* parasite. *Biochem Cell Biol* 2008, **86**:331-44.
62. Lu HL, Vinson SB, Pietrantoni PV: Oocyte membrane localization of vitellogenin receptor coincides with queen flying age, and receptor silencing by RNAi disrupts egg formation in fire ant virgin queens. *FEBS J* 2009, **276**:3110-23.
63. Amdam GV, Norberg K, Page RE Jr, Erber J, Scheiner R: Downregulation of vitellogenin gene activity increases the gustatory responsiveness of honey bee workers (*Apis mellifera*). *Behav Brain Res* 2006, **169**:201-5.
64. Marco Antonio DS, Guidugli-Lazzarini KR, do Nascimento AM, Simões ZL, Hartfelder K: RNAi-mediated silencing of vitellogenin gene function turns honeybee (*Apis mellifera*) workers into extremely precocious foragers. *Naturwissenschaften* 2008, **95**:953-61.
65. Sorokin AV, Kim ER, Ovchinnikov LP: Proteasome system of protein degradation and processing. *Biochem (Mosc)* 2009, **74**:1411-42.
66. Steller H: Regulation of apoptosis in *Drosophila*. *Cell Death Differ* 2008, **15**:1132-1138.
67. de la Fuente J, Blouin EF, Manzano-Roman R, Naranjo V, Almazán C, Perez de la Lastra JM, Zivkovic Z, Jongejan F, Kocan KM: Functional genomic studies of tick cells in response to infection with the cattle pathogen, *Anaplasma marginale*. *Genomics* 2007, **90**:712-722.
68. Hajdusek O, Sojka D, Kopecek P, Buresova V, Franta Z, Sauman I, Winzerling J, Grubhoffer L: Knockdown of proteins involved in iron metabolism limits tick reproduction and development. *Proc Natl Acad Sci USA* 2009, **106**:1033-8.
69. Beyenbach KW, Wiczkorek H: The v-type H<sup>+</sup> ATPase: molecular structure and function, physiological roles and regulation. *J Exp Biol* 2006, **209**:577-589.
70. de Vito P: The sodium/hydrogen exchanger: A possible mediator of immunity. *Cell Immunol* 2006, **240**:69-85.
71. Whyard S, Singh AD, Wong S: Ingested double-stranded RNAs can act as species-specific insecticides. *Insect Biochem Mol Biol* 2009, **39**:824-32.
72. Hao L, Sakurai A, Watanabe T, Sorensen E, Nidom CA, Newton MA, Ahlquist P, Kawaoka Y: *Drosophila* RNAi screen identifies host genes important for influenza virus replication. *Nature* 2008, **454**:890-893.
73. Kocan KM, Zivkovic Z, Blouin EF, Naranjo V, Almazán C, Mitra R, de la Fuente J: Silencing of genes involved in *Anaplasma marginale*-tick interactions affects the pathogen developmental cycle in *Dermacentor variabilis*. *BMC Development Biol* 2009, **9**:42.
74. Lundgren J, Masson P, Realini CA, Young P: Use of RNA interference and complementation to study the function of the *Drosophila* and human 26S proteasome subunit S13. *Mol Cell Biol* 2003, **23**:5320-30.
75. Wójcik C, DeMartino GN: Analysis of *Drosophila* 26S proteasome using RNA interference. *J Biol Chem* 2002, **277**:6188-97.
76. Hornok S, Földvári G, Elek V, Naranjo V, Farkas R, de la Fuente J: Molecular identification of *Anaplasma marginale* and rickettsial endosymbionts in blood-sucking flies (Diptera: Tabanidae, Muscidae) and hard ticks (Acari: Ixodidae). *Vet Parasitol* 2008, **154**:354-9.
77. Zhang B, McGraw E, Floate KD, James P, Jorgensen W, Rothwell J: *Wolbachia* infection in Australasian and North American populations of *Haematobia irritans* (Diptera: Muscidae). *Vet Parasitol* 2009, **162**:350-3.
78. Habayeb MS, Ekengren SK, Hultmark D: Nora virus, a persistent virus in *Drosophila*, defines a new picorna-like virus family. *J Gen Virol* 2006, **87**:3045-3051.



79. Eleftherianos I, Millichap PJ, French-Constant RH, Reynolds SE: **RNAi suppression of recognition protein mediated immune responses in the tobacco hornworm *Manduca sexta* causes increased susceptibility to the insect pathogen *Photobacterium*.** *Dev Comp Immunol* 2006, **30**:1099-107.
80. Ayres JS, Freitag N, Schneider DS: **Identification of *Drosophila* mutants altering defense of and endurance to *Listeria monocytogenes* infection.** *Genetics* 2008, **178**:1807-15.
81. Han-Ching Wang K, Tseng CW, Lin HY, Chen IT, Chen YH, Chen YM, Chen TY, Yang HL: **RNAi knock-down of the *Litopenaeus vannamei* Toll gene (LvToll) significantly increases mortality and reduces bacterial clearance after challenge with *Vibrio harveyi*.** *Dev Comp Immunol* 2010, **34**:49-58.
82. Schetinger MR, Morsch VM, Bonan CD, Wyse AT: **NTPDase and 5'-nucleotidase activities in physiological and disease conditions: new perspectives for human health.** *Biofactors* 2007, **31**:77-98.
83. Liyou N, Hamilton S, McKenna R, Elvin C, Willadsen P: **Localisation and functional studies on the 5'-nucleotidase of the cattle tick *Boophilus microplus*.** *Exp Appl Acarol* 2000, **24**:235-46.
84. Faudry E, Lozzi SP, Santana JM, D'Souza-Ault M, Kieffer S, Felix CR, Ricart CA, Sousa MV, Vernet T, Teixeira AR: ***Triatoma infestans* apyrases belong to the 5'-nucleotidase family.** *J Biol Chem* 2004, **279**:19607-13.
85. Valenzuela JG, Garfield M, Rowton ED, Pham VM: **Identification of the most abundant secreted proteins from the salivary glands of the sand fly *Lutzomyia longipalpis*, vector of *Leishmania chagasi*.** *J Exp Biol* 2004, **207**:3717-29.
86. Van Den Abbeele J, Caljon G, Dierick JF, Moens L, De Ridder K, Coosemans M: **The *Glossina morsitans* tsetse fly saliva: general characteristics and identification of novel salivary proteins.** *Insect Biochem Mol Biol* 2007, **37**:1075-85.
87. Francischetti IM, Valenzuela JG, Pham VM, Garfield MK, Ribeiro JM: **Toward a catalog for the transcripts and proteins (sialome) from the salivary gland of the malaria vector *Anopheles gambiae*.** *J Exp Biol* 2002, **205**:2429-51.
88. Francischetti IM, Meng Z, Mans BJ, Gudderra N, Hall M, Veenstra TD, Pham VM, Kotsyfakis M, Ribeiro JM: **An insight into the salivary transcriptome and proteome of the soft tick and vector of epizootic bovine abortion, *Ornithodoros coriaceus*.** *J Proteomics* 2008, **71**:493-512.
89. Schmidt CD, Harris RL, Hoffman RA: **Mass rearing of the horn fly, *Haematobia irritans* (Diptera: Muscidae), in the laboratory.** *Ann Entomol Soc Am* 1967, **60**:508-510.
90. Ririe KM, Rasmussen RP, Wittwer CT: **Product differentiation by analysis of DNA melting curves during the polymerase chain reaction.** *Anal Biochem* 1997, **245**:154-160.
91. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta CT) Method.** *Methods* 2001, **25**:402-408.

doi:10.1186/1471-2164-12-105

**Cite this article as:** Torres et al.: Functional genomics of the horn fly, *Haematobia irritans* (Linnaeus, 1758). *BMC Genomics* 2011 **12**:105.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

